


# A GENERALIZED INFLATION MODEL WITH COSMIC GRAVITATIONAL WAVES

V.N. LUKASH<sup>1</sup>, E.V. MIKHEEVA<sup>1</sup>, V. MÜLLER<sup>2</sup>, A.M. MALINOVSKY<sup>1</sup>

<sup>1</sup> *Astro Space Center of Lebedev Physical Institute,  
84/32 Profsoyuznaya, Moscow 117810, Russia*

<sup>2</sup> *Astrophysikalisches Institute Potsdam, An der Sternwarte 16,  
Potsdam, D-14482 Germany*



We propose a  $\Lambda$ -inflation model which explains a large fraction of the COBE signal by cosmic gravitational waves. The primordial density perturbations fulfil both the constraints of large-scale microwave background and galaxy cluster normalization. The model is tested against the galaxy cluster power spectrum and the high-multipole angular CMB anisotropy.

## 1 Introduction

The observational reconstruction of the *cosmological density perturbation* (CDP) spectrum is a key problem of the modern cosmology. It provides a dramatic challenge after detecting the primordial CMB anisotropy as the signal found by DMR COBE at  $10^0$  has appeared to be few times higher than the expected value of  $\Delta T/T$  in the most simple and best developed cosmological standard CDM model.

During recent years there were many proposals to improve  $\Lambda$ CDM (in the simplest term, to remove the discrepancy between the CDP amplitudes at  $8h^{-1}\text{Mpc}$  as determined by galaxy clusters, and at large scales,  $\sim 1000h^{-1}\text{Mpc}$ , according to  $\Delta T/T$ ) by adding hot dark matter, a  $\Lambda$ -term, or considering non-flat primordial CDP spectra. Below, we present another, presumably more natural way to solve the  $\Lambda$ CDM problem based on taking into account a possible contribution of *cosmic gravitational waves* (CGWs) into the large-scale CMB anisotropy; we will also try to preserve the original near-scale-invariant CDP spectrum. Thus, the problem is reduced to the construction of a simple inflation producing near *Harrison-Zel'dovich* (HZ) spectrum of CDPs ( $n_S \simeq 1$ ) and a considerable contribution of CGWs into the large-scale  $\Delta T/T$ .

A simple model of such kind is  $\Lambda$ -inflation, an inflationary model with an effective metastable  $\Lambda$ -term<sup>1, 2</sup>. This model produces both S (CDPs) and T (CGWs) modes which have a non-power-law spectra, with a shallow minimum in the CDP spectrum, located at a scale  $k_{cr}$  (there the  $\Lambda$ -term and the scalar field have equal energies while slowly-rolling at inflation) where the S-slope is exact HZ locally. The S-spectrum is 'red' for  $k < k_{cr}$ , and 'blue' for  $k > k_{cr}$ ; around the  $k_{cr}$  scale T/S is close to its maximum, it is of the order unity depending on the model parameters.

## 2 $\Lambda$ -inflation with self-interaction

Let us consider a general potential of  $\Lambda$ -inflation driven by a single scalar field  $\varphi$ :

$$V(\varphi) = V_0 + \sum_{\kappa=2}^{\kappa_{max}} \frac{\lambda_{\kappa}}{\kappa} \varphi^{\kappa}, \quad (1)$$

where  $V_0 > 0$  and  $\lambda_{\kappa}$  are constants,  $\kappa = 2, 3, 4, \dots$ . In the case of massive inflaton ( $\kappa = \kappa_{max} = 2$ ) T/S can be larger than unity only when the CDP spectrum slope in the 'blue' asymptote is very steep,  $n_S^{blue} > 1.8$ . To avoid such a strong spectral bend on short scales ( $k > k_{cr}$ ) we choose here another simple version of  $\Lambda$ -inflation – the case with self-interaction:  $\kappa = \kappa_{max} = 4$ ,  $\lambda_4 \equiv \lambda > 0$ ; this model is called  $\Lambda\lambda$ -inflation.

The scalar field  $\varphi$  drives an inflationary evolution if  $\gamma \equiv -\dot{H}/H^2 < 1$ , where  $H \simeq \sqrt{V/3}$  (we assume  $8\pi G = c = \hbar = 1$  and  $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$ ). This condition holds true for all values of  $\varphi$  if

$$c \equiv \frac{1}{4} \varphi_{cr}^2 = \frac{1}{2} \sqrt{\frac{V_0}{\lambda}} > 1, \quad (2)$$

which we imply hereafter. The fundamental gravitation perturbation spectra  $q_k$  and  $h_k$  generated in  $\Lambda\lambda$ -inflation in S and T modes, respectively, are as follows<sup>3, 4, 2</sup>:

$$q_k = \frac{H}{2\pi\sqrt{2\gamma}} = \frac{\sqrt{2\lambda/3}}{\pi} (c^2 + x^2)^{3/4}, \quad h_k = \frac{H}{\pi\sqrt{2}} = \frac{2c\sqrt{\lambda/3}}{\pi} \left(1 + \frac{x}{\sqrt{c^2 + x^2}}\right)^{-1/2}, \quad (3)$$

where

$$x = \ln \left[ \frac{k}{k_{cr}} \left(1 + \left(\frac{x}{c}\right)^2\right)^{1/4} \left(1 + \frac{x}{\sqrt{c^2 + x^2}}\right)^{2/3} \right] \simeq \ln(k/k_{cr}). \quad (4)$$

The dimensionless spectrum of density perturbations depends on a transfer function  $T(k)$ :

$$\Delta_k = 3.6 \times 10^6 \left(\frac{k}{h}\right)^2 q_k T(k). \quad (5)$$

## 3 CDM cosmology from $\Lambda\lambda$ -inflation

Let us consider the CDP spectrum with CDM transfer function, normalized both by  $\Delta T/T|_{10^0}$  (including the contribution from CGWs) and the galaxy cluster abundance at  $z = 0$ , to find the family of the most realistic  $q$ -spectra produced in  $\Lambda\lambda$ -inflation.

In total, we have three parameters entering the function  $q_k$ :  $\lambda$ ,  $c$  and  $k_{cr}$ . Constraining them by two observational tests we are actually left with only one free parameter (say,  $k_{cr}$ ) which may be restricted elsewhere by other observations.

To demonstrate explicitly how the three parameters are mutually related, we first employ a simple analytical estimates for the  $\sigma_8$  and  $\Delta T/T$  tests to derive the key equation relating  $c$  and  $k_{cr}$ , and then solve it explicitly to obtain the range of interesting physical parameters.

Instead of taking the  $\sigma$ -integral numerically we may estimate the spectrum amplitude on cluster scale ( $k = k_1 \simeq 0.3h/\text{Mpc}$ ):

$$q_{k_1} \simeq 4.5 \times 10^{-7} \frac{h^2 \sigma_8}{k_1^2 T(k_1)}. \quad (6)$$

On the other hand, the spectrum amplitude on large scale ( $k_2 = k_{\text{COBE}} \simeq 10^{-3}h/\text{Mpc}$ ) can be taken from  $\Delta T/T$  due to the Sachs-Wolfe effect<sup>5</sup>:

$$\langle \left( \frac{\Delta T}{T} \right)^2 \rangle_{10^0} = S + T \simeq 1.1 \times 10^{-10}, \quad S = 0.04 \langle q^2 \rangle_{10^0} \simeq 0.06 q_{k_2}^2. \quad (7)$$

The relation between the variance of the  $q$  potential averaged in  $10^0$ -angular-scale and the power spectrum at COBE scale, involves a factor of the effective interval of spectral wavelengths proportional to  $\ln \left( \frac{k_2}{k_{\text{hor}}} \right) \simeq 1.6$ . To estimate  $T/S$ , we use the following approximation formula at  $x_2 = x_{\text{COBE}}$ :

$$\frac{T}{S} \simeq -6n_T = \frac{6}{\sqrt{c^2 + x_2^2}} \left( 1 - \frac{x_2}{\sqrt{c^2 + x_2^2}} \right). \quad (8)$$

Evidently, both normalizations, (6) and (7), determine essentially the corresponding  $q_k$  amplitudes at the locations of cluster ( $k_1$ ) and COBE ( $k_2$ ) scales. Taking their ratio we get the key equation relating  $c$  and  $k_{cr}$  (see eqs.(3),(4),(8)):

$$\left( \frac{q_{k_1}}{q_{k_2}} \right)^2 \simeq D \left( 1 + \frac{T}{S} \right), \quad (9)$$

Eq.(9) has a clear physical meaning: the ratio of the S-spectral powers at cluster and COBE scales is proportional to  $\sigma_8^2$  and inversely proportional to the fraction of the scalar mode contributing to the large-scale temperature anisotropy variance,  $S/(S+T)$ . It provides quite a general constraint on the fundamental inflation spectra in a wide set of dark matter models using only two basic measurement (the cluster abundance and large scale  $\Delta T/T$ ). The DM information is contained in the D-coefficient which can be calculated using the same equation (9) for a simple inflationary spectrum (preserving the given DM model). For CDM with  $h = 0.5$  we have:

$$D \simeq \frac{0.6\sigma_8^2}{1 - 3.1\Omega_b}, \quad \Omega_b < 0.2. \quad (10)$$

The solution of eq.(9) has been obtained in the plane  $x_2 - c$  numerically. For the whole range  $0.1 < D < 0.5$ , it can be analytically approximated with a precision better than 10% as follows:

$$\ln^2 \left( \frac{k_0}{k_{cr}} \right) \simeq E (c_0 - c) (c + c_1), \quad 2 < c \leq c_0. \quad (11)$$

Notice there exists no solution of eq.(9) for  $c > c_0$ . We have found the following best fit coefficients  $E$ ,  $k_0[h/\text{Mpc}]$  and  $c_{0,1}$ :

$$E \simeq 1, \quad \ln k_0 \simeq 49D^2 + 1.3, \quad c_0 \simeq 61D^2 + 6.2, \quad c_1 \simeq 44D^2 + 4.0.$$

The tensor-mode-contribution is approximated similarly ( $k_{cr}$  is measured in  $[h/\text{Mpc}]$ ):

$$\frac{T}{S} \simeq \frac{2.53 - 4.3D}{(\ln k_{cr} + 4.65)^{2/3}} + \frac{1}{3}. \quad (12)$$

## 4 Discussion

We have presented a new inflationary model predicting a near scale-invariant spectrum of density perturbations and large amount of CGWs. Our model is consistent with COBE  $\Delta T/T$  and cluster abundance data. The perturbation spectra depend on one free scale-parameter,  $k_{cr}$ , which can be found in further analysis by fitting other observational data. At the location of  $k_{cr}$ , the CDP spectrum transfers smoothly from the red ( $k < k_{cr}$ ) to the blue ( $k > k_{cr}$ ) parts.

Today we seriously discuss a nearly flat shape of the dimensionless CDP spectrum within the scale range encompassing clusters and superclusters,

$$\Delta_k^2 \sim k^{(0.9^{+0.2})}, \quad k \in (0.04, 0.2)h \text{ Mpc}^{-1}, \quad (13)$$

(with a break towards the HZ slope on higher scales) which stays in obvious disagreement with the  $\Lambda$ CDM prediction. The arguments supporting eq.(13) came from the analysis of large-scale galaxy distribution<sup>6</sup> and the discovery of large quasar groups<sup>7, 8</sup>, a higher statistical support was brought by recent measurements of the galaxy cluster power spectrum<sup>9, 10</sup>.

A possible explanation of eq.(13) could be a fundamental red power spectrum established on large scales, then the transition to the spectrum (13) at  $\sim 100\text{Mpc}/h$  would be much easier understood with help of a traditional modification of the transfer function  $T(k)$  (e.g. for mixed hot+cold dark matter). The redness may be not too high, remaining in the range (0.9, 1). A way to enhance the power spectrum at Mpc scale could be the identification of  $k_{cr}$  within a cluster scale ( $k_{cr} \sim k_1$ ).

Notice that one of the problems for the matter-dominated models is a low number of  $\sigma_8$ : if  $\sigma_8 < 0.6$ , then the first acoustic peak in  $\Delta T/T$  cannot be as high as  $\gtrsim 70 \mu\text{K}$ .

### Acknowledgments

The work was partly supported by German Scientific Foundation (DFG-436 RUS 113/357/0) and INTAS grant (97-1192). V.N.L. and E.V.M. are grateful to the Organizing Committee for the hospitality.

1. V.N. Lukash, E.V. Mikheeva, *Gravitation & Cosmology* **2**, 247 (1996).
2. V.N. Lukash, E.V. Mikheeva, *Int.J.Mod.Phys.A*, in press.
3. V.N. Lukash, *Zh. Eksp. Teor. Fiz.* **79**, 1601 (1980).
4. V.N. Lukash in *Cosmology: The Physics of the Universe*, eds. B A Robson et al. (World Scientific, Singapore, 1996).
5. R.K. Sachs, A.M. Wolfe, *ApJ* **147**, 73 (1967).
6. L. Guzzo et al., *ApJ* **382**, L5 (1991).
7. B.V. Komberg, V.N. Lukash, *MNRAS* **269**, 277 (1994).
8. A.V. Kravtsov, B.V. Komberg, V.N. Lukash, *A&A* **282**, 713 (1996).
9. H. Tadros, G. Efstathiou, C. Dalton, *MNRAS* **296**, 995 (1998).
10. J. Retzlaff et al., *NewA* **3**, 631 (1998).